



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

GENERAL NOTES.

The Black Tern at Philadelphia, Pa.— On October 20, 1906, returning from an unsuccessful ducking trip behind Petty's Island, in the Delaware River, opposite Philadelphia, we (my brother and myself) observed a Black Tern (*Hydrochelidon nigra surinamensis*) which was flushed from the water about two hundred yards off Williams Street wharf, by a passing tug. It flew slowly up the river, keeping close to the shore, but beyond gunshot. It was plainly visible to us, as we were not over a hundred yards distance from it when it took wing, and the black color and white under tail coverts precluded any possibility of doubt as to its identity.

The Black Tern is given in Stone's 'Birds of Eastern Pennsylvania and New Jersey,' (p. 32) as a "rare or irregular transient" in this vicinity, and in the same book (p. 48) he says, "stragglers have been reported from the Delaware and Susquehanna Rivers," but I can find no records of its occurrence on the former river. Regarding this species in New Jersey, Mr. Stone says (p. 48 of his book), "transient, occurring during the fall migration on the New Jersey coast"; and from this remark I infer that the bird seen by us was one of these transients, driven inland by the stormy weather and northeast winds that prevailed for about a week previous to our observation. Its presence here cannot be attributed to any other cause, which is undoubtedly the true reason of its occurrence.

The Greater Yellow legs (*Totanus melanoleucus*) occurs here only after northeast storms, which also occasionally drive inland other rare species of water birds from the New Jersey coast. The Common Terns (*Sterna hirundo*) are sometimes abundant on the Delaware River above Philadelphia during northeast storms and always disappear after their abatement.

Mr. Stone is correct in considering the Black Tern a "rare or irregular transient" in this vicinity, for it is of such extremely rare occurrence that there is no record of its capture during recent years on the Delaware.—
RICHARD F. MILLER, Philadelphia Pa.

Ross's Snow Goose in Colorado.— On December 23, 1906, there was shot on the Kennicott Club Lake, $3\frac{1}{2}$ miles east of Longmont, Colo., a male Ross's Snow Goose (*Chen rossii*). The bird was killed by Capt. Eli, U. S. A., presented by him to Mr. Mart H. Watrous, mounted by Mr. Rudolph Borchardt, and later generously presented by Mr. Watrous to the Colorado Museum of Natural History, Denver, where it may now be seen on exhibition. It was associating with a flock of Mallards (*Anas boschas*), with which it had also been seen by the lake keeper the day before. Two or three days previous to this time the keeper's wife had seen a "flock of white geese about the same size as this one," but whether they were Ross's Geese or not can only be conjectured. I have examined the bird very carefully, comparing it with a specimen from my own col-